

Learning Objective: To review probability techniques including venn diagrams and two-way tables.

### Conditional probability using Venn diagrams

**Conditional probability** refers to the chances that some outcome occurs given that another event has also occurred.

The notation for conditional probability is P(B|A). This reads 'the probability of B given A'.

This is calculated as:

*P* (*B*\*A*) = number of elements in *B* and *A* ÷ number of elements in *A* 

### Example:

The following Venn diagram shows the number of students in year 10, where:

C = the number of students in the Chess club. M = the number of students in the Maths club.

One student is chosen at random. Find the probability that the student is in the Chess club, given that they are also in the Maths club.

 $P(C \mid M) = \frac{number of elements in C and M}{number of elements in M}$ 

<u>7</u> 13

### The tree diagram below shows the outcomes at each stage of an experiment tossing a coin three times.

An ordinary die with faces labelled 1 to 6 has had one corner sliced off, thus creating a seventh face labelled 7. The results of throwing the die 60 times are shown below:





Find the probability, in simplified fraction form, that the number rolled is:

A prime number.	
A factor of 30.	
An even number.	

# A medical researcher studies 1000 people to see whether they contracted an illness and whether they were vaccinated against it. Find the probability:

Suffered illness Vaccinated	A randomly selected person avoided the illness.
94 3 328	A randomly selected person was vaccinated, find the probability that they got the illness.
575	A randomly selected person did not get ill, find the probability that they were vaccinated.



## MATHEMATICS: ANSWER SHEET

Learning Objective: To review probability techniques including venn diagrams and two-way tables.

### Conditional probability using Venn diagrams

**Conditional probability** refers to the chances that some outcome occurs given that another event has also occurred.

The notation for conditional probability is P(B|A). This reads 'the probability of B given A'.

This is calculated as:

*P* (B\A) = number of elements in B and A  $\div$  number of elements in A

### Example:

The following Venn diagram shows the number of students in year 10, where:

C = the number of students in the Chess club. M = the number of students in the Maths club.

 $= \frac{7}{13}$ 

One student is chosen at random. Find the probability that the student is in the Chess club, given that they are also in the Maths club.

 $P(C \setminus M) = \frac{number of elements in C and M}{number of elements in M}$ 

#### The tree diagram below shows the outcomes at each stage of an experiment tossing a coin three times.

An ordinary die with faces labelled 1 to 6 has had one corner sliced off, thus creating a seventh face labelled 7. The results of throwing the die 60 times are shown below:





Find the probability, in simplified fraction form, that the number rolled is:

A prime number.	P (prime) = $\frac{8+9+11+6}{60}$ = 34/60 = 17/30
A factor of 30.	P (factor of 30) = $\frac{6+8+9+11}{60}$ = 34 / 60 = 17 / 30
An even number.	P (even) = $\frac{8 + 10 + 10}{60}$ = 28 / 60 = 7 / 15

# A medical researcher studies 1000 people to see whether they contracted an illness and whether they were vaccinated against it. Find the probability:

Suffered illness Vaccinated 94 3 328 575	A randomly selected person avoided the illness.	P ('I) = $\frac{328 + 575}{94 + 3 + 328 + 575}$ = $\frac{903}{1000}$
	A randomly selected person was vaccinated, find the probability that they got the illness.	P (I\V) = $\frac{3}{328 + 3}$ = $\frac{3}{331}$
	A randomly selected person did not get ill, find the probability that they were vaccinated.	P (V\l') = $\frac{328}{328 + 575}$ = $\frac{328}{903}$